

# Rotation and Spin in Physics\*

R. F. O'Connell

*Department of Physics and Astronomy,  
Louisiana State University, Baton Rouge, LA 70803-4001*

(Dated: August 25, 2010)

## Abstract

We delineate the role of rotation and spin in physics, discussing in order Newtonian classical physics, special relativity, quantum mechanics, quantum electrodynamics and general relativity. In the latter case, we discuss the generalization of the Kepler formula to post-Newtonian order ( $c^{-2}$ ) including spin effects and two-body effects. Experiments which verify the theoretical results for general relativistic spin-orbit effects are discussed as well as efforts being made to verify the spin-spin effects.

---

\* This paper is dedicated to the memory of Professor J. A. Wheeler

## I. INTRODUCTION

This article is essentially a companion to an article which I recently prepared for the 2007 Varenna summer school [1]. The latter deals in detail with spin effects in general relativity at the post-Newtonian level (order  $c^{-2}$ ) with particular emphasis on the theory, and related experiments, of the two-body Kepler problem extended to include spin. Here, we use "spin" in the generic sense of meaning "internal spin" in the case of an elementary particle and "rotation" in the case of a macroscopic body.

My work with Barker on the general relativistic theory [2–4] brought to the fore some interesting conceptual matters dealing with the relation between velocity and momentum (especially for particles with spin), the non-uniqueness of spin supplementary conditions and the choice of coordinates even at the classical special relativistic level, the fact that a spinning particle necessarily has a minimum radius, the corresponding concepts in quantum theory (relating to such topics as the Foldy-Wouthuysen transformation and position operators) and the fact that the spin effects in quantum electrodynamics (obtained from one-photon exchange) have their analogy in general relativity (obtained from one-graviton exchange or purely classical calculations) to the extent that except for (important) numerical factors, the latter results may be obtained from the former by simply letting  $e^2 \rightarrow Gm_1m_2$ .

Thus, we are motivated to present in a systematic way the role of rotation and spin in physics generally. So, we discuss, in order, Newtonian classical physics, special relativity, quantum mechanics, quantum electrodynamics and general relativity, in sections 2 to 6, respectively. Our conclusions are presented in Section 7. The closest discussion of this nature in the literature is the book by Corben [5]. However, whereas his list of references proved a useful resource, his emphasis is on elementary particles and he does not consider general relativity, which is our emphasis.

## II. NEWTONIAN CLASSICAL PHYSICS

Newton's second law is generally written in the form

$$\vec{F} = m \frac{d\vec{v}}{dt}, \tag{2.1}$$

where  $\vec{v}$  is the velocity. Next, the momentum  $\vec{p}$  of a particle is essentially defined as  $m\vec{v}$  so that we have

$$\vec{F} = \frac{d\vec{p}}{dt}. \quad (2.2)$$

As French has emphasized [6], Newton in the *Principia* did not write (2.1) but in essence, wrote (2.2). Also, only in certain frames of reference, so-called inertial frames, are Newton's laws valid. All frames, moving uniformly in a straight line relative to a particular inertial frame, constitute an infinity of inertial frames where "the properties of space and time are the same, and the laws of mechanics are the same" [7], and such "frame(s) of reference can always be chosen in which space is homogenous and isotropic and time is homogeneous" [7]. This leads us to a new concept which assigns a broader independent meaning to momentum. It follows from the introduction of the Lagrangian  $L(q, \dot{q}, t)$  of a system, where  $q$  refers to position and  $\dot{q} = (dq/dt)$  is the velocity. As a result, the associated Euler-Lagrange equations for a closed system lead to conservation laws of energy (resulting from the homogeneity of time), momentum (from the homogeneity of space) and angular momentum (from the isotropy of space) where, in particular, the momentum is referred to as the canonical momentum  $\vec{P}$ , as distinct from the mechanical momentum  $\vec{p} = m\vec{v}$ , where

$$\vec{P} = \frac{\partial L}{\partial \vec{v}}. \quad (2.3)$$

In general,  $\vec{P}$  and  $\vec{p}$  are not equal, an example being the case of a particle with charge  $q$  in an electromagnetic field, where the force is velocity dependent, in which case

$$\vec{P} = \vec{p} + \frac{q\vec{A}}{c}. \quad (2.4)$$

Thus, whereas  $\vec{P}$  is conserved,  $\vec{p}$  is not conserved in general. Moreover, in the passage to quantum mechanics, it is the canonical variables which are of importance in forming operators. In addition, it is  $\vec{P}$  that enters into the conserved angular momentum  $\vec{L} = \vec{r} \times \vec{P}$ . Of course, in this case, the reason why  $\vec{P}$  and  $\vec{v}$  are not proportional is due to the presence of an external field. However, in special relativity, we shall see that, even for a free particle, even  $\vec{p}$  and  $m\vec{v}$  are, in general, not equal to each other.

Turning to the rotation of macroscopic bodies, we define  $\vec{S}$  as the spin or intrinsic angular momentum. In general,

$$S_i = I_{ij}\omega_j \quad (i, j = 1, 2, 3), \quad (2.5)$$

where  $\vec{\omega}$  is the angular velocity and  $I_{ij}$  are the components of the inertia tensor.

Based on observations, the earth is an inertial frame if one neglects its rotation. However, taking the "fixed stars" as an inertial system with "space axes", the earth rotates with an angular velocity  $\vec{\omega}$  relative to these axis, giving rise to Coriolis and centrifugal forces. Thus, following Goldstein [8] and writing

$$\vec{v}_s = \vec{v}_v + \vec{\omega} \times \vec{r}, \quad (2.6)$$

where  $\vec{v}_s$  and  $\vec{v}_r$  are the velocities of a particle relative to the space and the earth's rotating set of axis, the equation

$$\vec{F}_s = ma_s, \quad (2.7)$$

in the space system translates to

$$\vec{F}_r = ma_r = ma_s - 2m(\vec{\omega} \times \vec{v}_r) - m\vec{\omega} \times (\vec{\omega} \times \vec{r}), \quad (2.8)$$

in the rotating system. The second and third terms are the familiar Coriolis and centrifugal forces, respectively. Of particular interest is a result, not found in [8] but derived in [7], that the momentum of a particle is the same in both frames of reference so that, in the rotating system,

$$\vec{p} = m\vec{v} + m(\vec{\omega} \times \vec{r}). \quad (2.9)$$

Thus, here we have a case where the momentum is different from  $m\vec{v}$  purely due to the non-inertial nature of the rotating system. We also remark that the angular momenta are also equal in both systems but the energy in the rotating system  $E_r = E_s - \vec{L} \cdot \vec{\omega}$  [7].

Finally, we remark that the earth is not a perfect sphere but is in fact an "asymmetric top", so that  $\vec{\omega}$ , the angular velocity of the earth, as distinct from  $\vec{S}$  does not remain fixed in space but, instead, it executes what is known as polhode motion [7, 8]. This force-free precession of the earth's axis is a key to the explanation of Chandler's precession of the earth's axis and is also proving to be a major bugbear in the analysis of the GP-B gyroscope experiment [9, 10].

### III. SPECIAL RELATIVITY

Just as the angular momentum  $\vec{L}$  generalizes to an anti-symmetric second rank tensor so does the 3-vector  $\vec{S}$  generalizes to  $S_{\alpha\beta} = -S_{\beta\alpha}$ . Another possibility is to define an axial 4-vector  $S^\mu$  which reduces to the 3-vector  $\vec{S}$  in the rest-frame of the particle. In fact, this can be achieved by defining

$$S_\alpha \equiv \frac{1}{2} \epsilon_{\alpha\beta\sigma\tau} S^{\beta\sigma} U^\tau, \quad (3.1)$$

where  $\epsilon_{\alpha\beta\sigma\tau}$  is the completely antisymmetric Levy-Civita tensor,  $U^\tau = (\gamma, \vec{v}/c)$  is the familiar 4-velocity, and  $S^\alpha = (0, \vec{S})$  in the rest frame where  $\vec{v} = 0$ . Hence, using the fact that  $\epsilon_{\alpha\beta\sigma\tau}$  is antisymmetric in  $\alpha$  and  $\tau$ , we obtain

$$U^\alpha S_\alpha = 0, \quad (3.2)$$

so that the 4-vectors  $U^\alpha$  and  $S_\alpha$  are not only orthogonal in the rest frame as constructed but are also orthogonal in all frames. In addition, using the properties of the Levi-Civita symbol (Ref. [5], p. 79) and  $U^\alpha U_\alpha = -1$ , (3.1) may be inverted to give

$$S^{\alpha\beta} = \epsilon^{\alpha\beta\sigma\tau} S_\sigma U^\tau. \quad (3.3)$$

Eq. (3.2) is a spin supplementary condition (SSC) which ensures that even when  $\vec{v}$  is non-zero,  $S^\mu$  has only three independent components and similarly for  $S^{\alpha\beta}$ . For a free particle (no external forces or torques)

$$\frac{dS^\mu}{dt} = 0, \quad (3.4)$$

but, nevertheless, as detailed in [5], the particle moves in a circle in a plane normal to  $\vec{S}$  with a radius

$$\vec{r} = -\frac{\vec{v} \times \vec{S}}{mc^2}. \quad (3.5)$$

Such a motion is reminiscent of Zitterbewegung in relativistic quantum mechanics [11], as we shall discuss below. We will also see how such a quantity as  $\vec{r}$  appears in the discussion of spin effects on orbital motion in post-Newtonian general relativistic theory. For now, we note that such a problem can be circumvented by choosing a different SSC based on the fact that there are essentially two basic rest systems for the particle, corresponding to either  $\vec{v} = 0$  or  $\vec{p} = 0$ . This is also connected to the fact that, as Moller has shown [12, 13], a spinning body has a minimum radius equal to  $|\vec{S}|/mc$ . As a result, the definition of a rest frame is related to a choice of SSC which, in turn, is related to the choice of a center-of-mass for the spinning particle [14]. Thus, switching from one SSC (or rest-frame) to another is exactly the same as shifting the center of mass by a Lorentz transformation.

Turning to the case where a particle is acted on by a force  $f^\mu$  without experiencing any torque, Weinberg [15] has shown that

$$\frac{dS^\alpha}{d\tau} = \left( S_\beta \frac{f^\beta}{m} \right) U^\alpha, \quad (3.6)$$

which leads to the famous Thomas precession [16, 17]. The usual discussion of Thomas precession is concerned with precession of a 3-vector, leading to a reduction by  $\frac{1}{2}$  in the spin-orbit energy of an atomic electron (taking the gyromagnetic ratio  $g$  to be 2). This analysis was generalized by Bargmann et al. [18] to obtain  $dS^\alpha/d\tau$  in its most simple form by assuming from the outset that the momentum and velocity of a spinning particle are proportional [5]. Thus, they were led to the so-called BMT equation for the classical relativistic equation of motion for spin in uniform or slowly varying external fields and its various applications. Details of this work are given in Jackson's well-known book [17], along with the derivation of Thomas's equation of motion of the spin vector from which Thomas precession emerges. Important applications of the BMT equation are listed in [5].

#### IV. QUANTUM MECHANICS

As detailed in all the textbooks, spin was introduced into non-relativistic quantum mechanics by Goudsmit and Uhlenbeck [19]. It should also be mentioned that the Wigner distribution, which represents an alternative formulation of quantum mechanics [20, 21] has also been extended to incorporate spin [22].

Relativistic quantum mechanics was initiated by Dirac's equation for the electron. However, the Dirac equation for a free particle, leads to the conclusion that the momentum and velocity are not simply related [11]. Thus, whereas the plane-wave solutions are eigenfunctions of the momentum operator, the velocity operator is not a constant of the motion with the implication that the free electron not only displays uniform rectilinear motion but also executes very rapid oscillations, called Zitterbewegung by Schrodinger who, as early as 1930, first discussed such motion. However, since we noted earlier that all classical spinning bodies have a minimum radius  $|\vec{S}|/mc$ , it is reasonable to assume that the electron (for which  $|\vec{S}| = \hbar/2$ ) cannot be localized to a size smaller than  $\lambda_c/2$ , where  $\lambda = \hbar/mc$  is the Compton wavelength. Hence, analogous to the Lorentz transformations which shift the center of mass of a classical particle with spin, for an electron we can change the coordinate operator by means of a unitary Foldy-Wouthuysen transformation [23, 24]. In this representation the velocity operator is proportional to the momentum operator but, as emphasized by Sakurai (p. 177 of [11]), "nonlocality — is the price we must pay." By contrast, "—a well-localized state contains, in general, plane-wave components of negative energy" [11]. In earlier work and in a similar vein, Newton and Wigner [25] created the most localized state possible for a free particle without spin, and also the corresponding position operator, by using only plane-wave components of positive energies (solutions of the Klein-Gordon equation) and concluded that it is not a delta function as in the nonrelativistic case; instead, the wave function goes as  $r^{-5/2}$  for small  $r$  and falls off exponentially for large  $r$  and, in general its characteristic extension in space is  $\approx \lambda_c$ . This work was then extended to particles with spin for which position operators and quantum states were also obtained. The results obtained are unique subject to the restriction that the orbital angular momentum of the localized state has the value  $\ell = 0$ . However, for spin 1/2 particles, it was shown by the present author and Wigner [26] that another choice of position operator is possible, given by

$$\vec{Q} = \vec{q} + (\vec{p} \times \vec{\sigma})/p^2, \quad (4.1)$$

where  $\vec{q}$  is the original position operator,  $\vec{Q}$  is the new position operator, and where the components of  $\vec{\sigma}$  denote the Pauli spin matrices. In addition, it was shown [27] that

$$\langle \vec{q}(t) \rangle = \langle \vec{q}(0) \rangle + t \langle \vec{v} \rangle, \quad (4.2)$$

where here the velocity operator is  $\vec{v} = (c^2\vec{p}/E)$ , where  $E$  is the total energy, including the rest mass. Thus, the movement of the mean position of a free particle obeys the classical Newtonian equation except for  $m \rightarrow (E/c^2)$ . Moreover, a similar result was obtained [26] for  $\langle \vec{Q}(t) \rangle$ . Even for the case of a spin zero particle, (4.2) is a non-trivial result because, as emphasized in [27], "the situation in quantum mechanics is not immediately obvious because although  $\vec{p}$  has a natural definition - it is the generator of spatial translations of the state vector - this is not so for  $\vec{v}$  - the "velocity" does depend on the definition of the position". Finally, we note that the choice of state vectors given by (4.1) and (4.2) leads to motion free from Zitterbewegung, just as we saw for the Foldy-Wouthuysen position operator.

## V. QUANTUM ELECTRODYNAMICS (QED)

The initial (semi-classical) treatment of particles and atoms with the electromagnetic field considered the latter to be classical. This was dramatically changed in 1927 when Dirac quantized the electromagnetic field [28]. Thus, for fields in general, the emphasis then switched from trying to construct relativistic potentials to the consideration of suitable Lagrangians and the use of Feynman diagrams,  $S$ -matrix theory [29] and other techniques. These are particularly useful in carrying out perturbation calculations. In contrast to nonrelativistic quantum theory (where potentials are used) in relativistic field theory, interactions are considered to arise from the exchange of quanta. Thus, in QED, the quanta are photons of spin 1 (whereas, as we shall discuss at length in the next section, gravitational interactions can be considered to be mediated by spin 2 gravitons). In essence, each diagram gives rise to a covariant matrix element whose Fourier transform "enables us to construct an effective three-dimensional potential (to be used in connection with the Schrodinger equation)"



(Ref. [11], p. 259). A detailed description of this technique is given in [30]; in particular, they consider the electromagnetic interaction of two particles with masses  $m$ , and  $m_2$  (such as an electron and a muon) for which they first calculate the particle interaction operator in the momentum representation. Next, by taking Fourier transforms, they obtain the corresponding operator in the coordinate representation (Ref. [30], p. 283, Eq. (83.15)); the latter contains terms of purely orbital origin as well as spin-orbit and spin-spin terms, and Darwin-like terms. As we shall see in the next section, an analogous method for the gravitational interaction of two spinning bodies leads to similar type terms.

## VI. GENERAL RELATIVITY

For a free particle with spin  $S^\alpha$ , in special relativity we had the result given by (3.4). The covariant generalization of this formula is [15]

$$\frac{DS^\mu}{D\tau} = 0, \quad (6.1)$$

where  $D$  denotes covariant differentiation. Also, in the presence of a gravitational field, (3.6) generalizes to [15, 31]

$$\frac{DS^\mu}{D\tau} = S_\nu \frac{DU^\nu}{D\tau} U^\mu \equiv S_\nu a^\nu U^\mu, \quad (6.2)$$

where  $a^\mu = \frac{f^\mu}{m}$  is the 4-acceleration. This is referred to as Fermi-Walker transport of the spin vector in its motion along a curve  $x^\alpha$  with tangent vector  $U^\alpha$ . In the special case where  $f^\mu = 0$  (free fall), the right-side of (6.2) is zero and the Fermi-Walker transport reduces to parallel transport along the geodesic, given by (6.1), which may be written explicitly as

$$\frac{dS_\mu}{d\tau} = \Gamma^\lambda_{\mu\nu} S_\lambda \frac{dx^\nu}{d\tau}, \quad (6.3)$$

where  $\Gamma^\lambda_{\mu\nu}$  is the Christoffel symbol. Next, Papapetrou [32, 33] considered the motion of a small mass with spin in a gravitational field in order to obtain the deviation from geodesic motion. This deviation is expressed in both orbital and spin equations of motion, given by

$$\frac{D}{Ds} \left( mu^\alpha + u_\beta \frac{DS^{\alpha\beta}}{Ds} \right) + \frac{1}{2} S^{\mu\nu} u^\sigma R_{\nu\sigma\mu}^\alpha = 0, \quad (6.4)$$

and

$$\frac{DS^{\alpha\beta}}{Ds} + u^\alpha u_\rho \frac{DS^{\beta\rho}}{Ds} - u^\beta u_\rho \frac{DS^{\alpha\rho}}{Ds} = 0, \quad (6.5)$$

respectively, where  $R_{\nu\sigma\mu}^\alpha$  is the Riemann curvature tensor and  $S^{\alpha\beta}$  is given by (3.3). This was the starting-point used by Schiff [34] in his derivation of the theory underlying the GP-B experiment [10], which involved calculations to post-Newtonian order. However, Barker and the present author [2] applied a completely different approach to the problem than the method used by Papapetrou and Schiff, obtaining results which agreed with those of Schiff for the spin precession but apparently different for the orbital equations of motion.

The method used in [2] was based on a quantum approach to gravitation developed by Gupta [35] and his collaborators [36]. In particular, the gravitational interaction between two spin 1/2 Dirac particles was calculated by means of the exchange of a spin 2 graviton, analogous to what we discussed in Section 5 in the case of the electromagnetic interaction of two charged spin 1/2 Dirac particles. In fact, comparing the results in the latter case for the interaction of, say, an electron and muon (Ref. [30], Eq. (83.15)) with these obtained in the former case [2], we were able to conclude that all such effects in QED have their analogy in general relativity and, except for (important) numerical factors, the latter results may be obtained from the former by simply letting  $e^2 \rightarrow Gm_1m_2$ . Based on the universality of gravitational interactions, the transition from the microscopic to the macroscopic is achieved by the simple replacement  $\frac{1}{2} \hbar \vec{\sigma} \rightarrow \vec{S}$  [2]. In a later publication [37], we traced the apparent difference between our results and those of Schiff for the orbital equations of motion to the use of two different coordinate vectors, reflecting the choice of two different SSC's which in turn imply different choices for the center-of-mass of the gyroscope and also imply different relationships between the momentum and velocity vectors. Thus, as outlined in [37], Corinaldesi-Papapetrou and Schiff used  $S^{i0} = 0$  as their SSC; Pirani used  $S^{\alpha\beta}U_\beta = 0$  whereas Moller, Tulczyjew and Dixon used  $S^{\alpha\beta}P_\beta = 0$ . In particular, the difference in the coordinate vectors used by all investigators is always of the order  $(\vec{v} \times \vec{s})/mc^2$ , the same quantity which arose in the discussion of anomalous motion in special relativity [see (3.5)]

and not unrelated to the choice of position operators in relativistic quantum theory [see (4.1)]. However, regardless of the choice of SSC, in all cases identical results are obtained for orbital precessions. We also remark that, in contrast to the orbital case, the form of the spin equations of motion is the same for all choices of SSC's; the reason is that spin contributions are always order  $c^{-2}$  compared to the lowest-order Newtonian term appearing in the equation of motion.

Turning to experimental tests, Schiff's 1960 paper stimulated the birth of the GP-B experiment. However, because results are only now emerging, the initial euphoria which this experiment generated has waned [38] due to the fact that other investigators have verified the existence of spin-orbit effects in gravitation but, at least, the GP-B initial report [10] claims increased accuracy for the results obtained for the same quantity. The other investigations (which are discussed in more detail in [1], where the emphasis is on the Kepler problem and its post-Newtonian generalization, including spin) include (a) lunar-laser-ranging measurements of the lunar perigee [39, 40] based on the fact that the earth-moon system is essentially a gyroscope in the field of the sun [41]; (b) the Ciufolini Lageos experiment [42], based on the determination of earth-satellite distances with a precision of a few  $mm$ ; (c) the binary pulsar PSRB1534+12 observations [43] based on a determination of time evolutions involving the spin direction.

The advantage of these experiments is that they are on-going (in contrast to the GP-B experiment) and thus improved accuracy is likely to be achieved. In particular, Ciufolini *et al.*, whose *Nature* paper [42] is the first very credible demonstration of gravitomagnetism, are making use of the continual improvement in accuracy of earth gravity models [44].

The observations in [43] while limited in precession also provide an initial test of a 2-body result for the spin-orbit precession [3, 4]. What we found is that, for the spin precession of body 1, the replacement in the one-body formula is  $m_2 \rightarrow m_2 + (\mu/3)$  and, in the case of the spin precession of body 2, the replacement is  $m_1 \rightarrow m_1 + (\mu/3)$ , where  $\mu$  is the reduced mass. This result has now been verified by at least 7 other authors, the simplest being a calculation which makes use of the one-body result, followed by a transformation to the center-of-mass of the two-body system [45]. A potentially more promising candidate is the double binary system [46] for which spin precession angles of  $4.8^\circ/\text{yr}$  (pulsar A) and  $5.1^\circ/\text{yr}$  (pulsar B) have been predicted, based on our theoretical results [3, 4].

Finally, we remark on what we consider to be the most theoretically appealing alternative

to Einstein's theory in which the spin of matter, as well as its mass, plays a dynamical role. This idea was initiated by Cartan [47] and extended by Sciama [48] and Kibble [49] using gauge theory. Utiyama initially investigated Einstein's theory and the Cartan theory within the framework of the homogeneous and inhomogeneous (Poincaré) groups, respectively [50]. For a review, we refer to [51]. We investigated possible additional spin interactions based on this so-called torsion theory and we found that spin-spin contact interactions, additional to these associated with Einstein's theory, occur [52, 53]. However, as it turns out, they do not contribute to the macroscopic experiments presently being carried out.

## VII. CONCLUSIONS

In surveying the role of rotation and spin over the broad range of physics, several important concepts were considered. In non-relativistic classical physics, we recalled that Newton's second law only holds in an inertial system (in which case the mechanical momentum  $\vec{p} = m\vec{v}$ ) but does not hold in a rotating system which is non-inertial. In the latter case,  $\vec{p}$  is not proportional to  $\vec{v}$ . Also, the introduction of a Lagrangian  $L$  for a system brings into consideration a new type of momentum, the canonical momentum  $\vec{P}$ , which is the derivative of  $L$  with respect to  $\vec{v}$  and which, in the case of external fields, is different than  $\vec{p}$ . The intrinsic angular momentum (spin)  $\vec{S}$  also finds a natural definition within the Newtonian framework. However, in special relativity, one must treat spin as either a 4-vector  $S^\mu$  or else as a second rank antisymmetric tensor and define a relationship between them which requires the introduction of a spin supplementary condition (SSC). This led to the realization that there are essentially two basic rest systems, corresponding to either  $\vec{v} = 0$  or  $\vec{p} = 0$ , with the possibility of considering a variety of choices for a SSC instead of the choice given in (3.2). As a result, different results for  $\vec{p}$  in terms of  $\vec{v}$  emerge and, concomitantly, this also implies that the position vector of even a microscopic spinning particle is not an observable in special relativity.

Next, turning to quantum mechanics, we encountered similar phenomena. In particular, Zitterbewegung was found to be a feature of a particular choice of coordinate operator associated with Dirac's formulation of relativistic electron theory. It could be removed by carrying out a unitary transformation to a different coordinate for which Zitterbewegung disappears and simple proportionality between  $\langle \vec{p} \rangle$  and  $\langle \vec{v} \rangle$  holds. However, even for a

free particle with or without spin, it was found necessary to choose a very specific position operator if one uses only plane-wave components of positive energy, in which case the particle is non-local and has a characteristic extension  $\approx \lambda_c$ . Extending consideration to the QED domain, we saw that the modus operandi is to consider the interactions as arising from the exchange of spin1 photons, leading to spin-orbit, spin-spin and other terms. A further extension of these ideas to the realm of general relativity called for the exchange of spin 2 gravitons. This led to a systematic derivation of spin-orbit and spin-spin terms, to order  $c^{-2}$ , for a two-body binary system. We discussed experiments which have already verified the spin-orbit effects. In fact, given the latter, the results for the spin-spin effects must conform to the requirement that the total angular momentum (orbital + spin of body 1 + spin of body 2) is conserved.

#### NOTE ADDED IN PROOF

Our result for spin-orbit precession in a 2-body system [3] has been verified, to an accuracy of 13%, by Breton *et al.* [54] for the precession of pulsar *B* in the double binary system. In addition, in [55], we presented, in Table 2, a comparison of numerical results for a variety of one-body and two-body systems.

- 
- [1] R.F. O’Connell, ”Gravito-Magnetism in one-body and two-body systems: Theory and Experiment”, in *Atom Optics and Space Physics*, Proc. of Course CLXVIII of the International School of Physics ”Enrico Fermi”, Varenna, Italy, ed. E. Arimondo, W. Ertmer, W. Schleich and E.M. Rasel (IOS, Amsterdam, 2009).
  - [2] B.M. Barker and R.F. O’Connell, ”Derivation of the Equations of Motion of a Gyroscope from the Quantum Theory of Gravitation”, *Phys. Rev. D* **2**, 1428 (1970).
  - [3] B.M. Barker and R.F. O’Connell, ”The Gravitational Two Body Problem with Arbitrary Masses, Spins, and Quadrupole Moments”, *Phys. Rev. D* **12**, 329 (1975).
  - [4] B.M. Barker and R.F. O’Connell, ”General Relativistic Effects in Binary Systems”, in *Physics and Astrophysics of Neutron Stars and Black Holes: Proceedings of Course 65 of the International School of Physics ”Enrico Fermi”*, Varenna, Italy, 1975, ed. R. Giacconi and R. Ruffini

- (North-Holland, 1976), p. 437.
- [5] H.C. Corben, "Classical and Quantum Theories of Spinning Particles", (Holden-Day, Inc., San Francisco, 1968).
  - [6] A. P. French, "Newtonian mechanics", (W.W. Norton, New York, 1971).
  - [7] L.D. Landau and E.M. Lifshitz, "Mechanics", 3rd ed. (Pergamon Press, Oxford, 1976).
  - [8] H. Goldstein, "Classical Mechanics", 2nd ed. (Addison-Wesley, Reading, MA, 1992).
  - [9] G. Brumfiel, "Gravity probe falters", *Nature* **444**, 978 (2006).
  - [10] Stanford, "GP-B Mission Status Update", Dec. 22, 2006.
  - [11] J.J. Sakurai, "Advanced Quantum Mechanics", (Addison-Wesley, Reading, MA, 1967).
  - [12] C. Moller, *Commun. Dublin Inst. Adv. Stud.*, **A5**, 1 (1949).
  - [13] C. Moller, "The Theory of Relativity", 2nd ed. (Oxford University Press, 1972), p. 176.
  - [14] B.M. Barker and R.F. O'Connell, "Nongeodesic Motion in General Relativity", *Gen. Relativ. Gravit.* **5**, 539 (1974).
  - [15] S. Weinberg, "Gravitation and Cosmology" (Wiley and Sons, New York, 1972), p. 124.
  - [16] L.H. Thomas, *Nature* **117**, 514 (1926).
  - [17] J.D. Jackson, "Classical Electrodynamics", 3rd ed. (Wiley, New York, 1998).
  - [18] V. Bargmann, L. Michel and V.L. Telegdi, "Precession of the Polarization of Particles Moving in a Homogeneous Electromagnetic Field", *Phys. Rev. Lett.* **2**, 435 (1959).
  - [19] G.E. Uhlenbeck and S. Goudsmit, *Nature* **117**, 264 (1926).
  - [20] M. Hillery, R.F. O'Connell, M. Scully, and E. Wigner, "Distribution Functions in Physics: Fundamentals", *Phys. Rep.* **106**, 121-167 (1984).
  - [21] R.F. O'Connell, "The Wigner Distribution" in *Compendium of Quantum Physics*, ed. by D. Greenberger, B. Falkenburg, K. Hentschel and F. Weinert (Springer, New York, 2009).
  - [22] R.F. O'Connell and E.P. Wigner, "Manifestations of Bose and Fermi Statistics on the Quantum Distribution Function for Systems of Spin Zero and Spin One-Half Particles", *Phys. Rev. A* **30**, 2613 (1984).
  - [23] L. Foldy and S.A. Wouthuysen, "On the Dirac Theory of Spin 1/2 Particles and Its Non-Relativistic Limit", *Phys. Rev.* **78**, 29 (1950).
  - [24] M.E. Rose, "Relativistic Electron Theory" (Wiley, New York, 1961).
  - [25] T.D. Newton and E.P. Wigner, "Localized States for Elementary Systems", *Rev. Mod. Phys.* **21**, 400 (1949).

- [26] R.F. O'Connell and E.P. Wigner, "Position Operators for Systems Exhibiting the Special Relativistic Relation Between Momentum and Velocity", *Phys. Lett.* **67A**, 319 (1978).
- [27] R.F. O'Connell and E.P. Wigner, "On the Relation Between Momentum and Velocity for Elementary Systems", *Phys. Lett.* **61A**, 353 (1977).
- [28] P.A.M. Dirac, *Proc. Roy. Soc. A* **114**, 243 (1927).
- [29] J.A. Wheeler, "On the Mathematical Description of Light Nuclei by the Method of Resonating Group Structure", *Phys. Rev.* **52**, 1107 (1937).
- [30] V.B. Berestetskii, E.M. Lifshitz, and L.P. Pitaevskii, *Relativistic Quantum Theory* (New York, N.Y., 1971), p. 280.
- [31] I. Ciufolini and J.A. Wheeler, "Gravitation and Inertia" (Princeton University Press, 1995).
- [32] A. Papapetrou, "Spinning Test-Particles in General Relativity I", *Proc. R. Soc. London* **A209**, 248 (1951).
- [33] E. Corinaldesi and A. Papapetrou, "Spinning Test-Particles in General Relativity II", *Proc. R. Soc. London* **A209**, 259 (1951).
- [34] L.I. Schiff, *Proc. Nat. Acad. Sci.* **46**, 871 (1960); *Proceedings of the Theory of Gravitation*, Jablonna, Poland, 1962, edited by L. Infeld (Paris, 1964), p. 71.
- [35] S.N. Gupta, "Einstein's and Other Theories of Gravitation", *Rev. Mod. Phys.* **29**, 334 (1957).
- [36] B.M. Barker, S.N. Gupta, and R.D. Haracz, "One-Graviton Exchange Interaction of Elementary Particles", *Phys. Rev.* **149**, 1027 (1966).
- [37] B.M. Barker and R.F. O'Connell, "Nongeodesic Motion in General Relativity", *Gen. Rel. Grav.* **5**, 539 (1974).
- [38] A. Lawler, "NASA Orders Make-or-Break Tests for Gravity Probe", *Science* **300**, 880 (2003).
- [39] J.G. Williams, X.X. Newhall, and J.O. Dickey, "Relativity parameters determined from lunar laser ranging", *Phys. Rev. D* **53**, 6730 (1996).
- [40] J.G. Williams, S.G. Turyshev and D.H. Boggs, "Progress in Lunar Laser Ranging Tests of Relativistic Gravity", *Phys. Rev. Lett.* **93**, 261101 (2004).
- [41] R.F. O'Connell, "A Note of Frame Dragging", *Class. Quantum Grav.* **22**, 3815 (2005).
- [42] I. Ciufolini and E.C. Pavlis, "A Confirmation of the General Relativistic Prediction of the Lense-Thirring Effect", *Nature* **431**, 958 (2004).
- [43] I.H. Stairs, S.E. Thoresett and Z. Arzoumanian, "Measurement of Gravitational Spin-Orbit Coupling in a Binary-Pulsar System", *Phys. Rev. Lett.* **93**, 141101 (2004).

- [44] I. Ciufolini, E.C. Pavlis and R. Peron, "Determination of frame-dragging using Earth gravity models from CHAMP and GRACE", *New Astronomy* **11**, 527 (2006).
- [45] L. Chan and R.F. O'Connell, "Two-Body Problems—A Unified, Classical, and Simple Treatment of Spin-Orbit Effects", *Phys. Rev. D* **15**, 3058 (1977).
- [46] M. Kramer et al., "Texts of General Relativity from Timing the Double Pulsar", *Science* **314**, 97 (2006).
- [47] E. Cartan, *C. R. Acad. Sci* **174**, 593 (1972), and *Ann. Ed. Norm. Sup.* **40**, 325 (1923), and **41**, 1 (1924), and **42**, 17 (1925).
- [48] D.W. Sciama, "The Physical Structure of General Relativity", *Rev. Mod. Phys.* **36**, 463, 1103 (1964).
- [49] T.W.B. Kibble, "Lorentz Invariance and the Gravitational Field", *J. Math. Phys. (N.Y.)* **2**, 212 (1961).
- [50] R. Utiyama, "Invariant Theoretical Interpretation of Interaction", *Phys. Rev.* **101**, 1597 (1956).
- [51] F.W. Hehl et al., "General Relativity with Spin and Torsion: Foundations and Prospects", *Rev. Mod. Phys.* **48**, 393 (1976).
- [52] R.F. O'Connell, "Contact Interactions in the Einstein and Einstein-Cartan-Sciama-Kibble (ECSK) Theories of Gravitation", *Phys. Rev. Lett.* **37**, 1653 (1976) and **38**, 298(E) (1977).
- [53] R.F. O'Connell, "Attractive Spin-Spin Contact Interactions in the ECSK Torsion Theory of Gravitation", *Phys. Rev. D* **16**, 1247 (1977).
- [54] R.P. Breton, V.M. Kaspi, M. Kramer, M.A. McLaughlin, M. Lyutikov, S.M. Ransom, I.H. Stairs, R.D. Ferdman, F. Camilo and A. Possenti, "Relativistic Spin Precession in the Double Pulsar," *Science* **321**, 104 (2008).
- [55] C. Feiler, M. Buser, E. Kajari, W.P. Schleich, E.M. Rasel and R.F. O'Connell, "New Frontiers at the Interface of General Relativity and Quantum Optics," *Space Sci. Rev.* **148**, 123-147 (2009).